

**Nuclear Data Section**  
**International Atomic Energy Agency**  
**P.O.Box 100, A-1400 Vienna, Austria**

**Memo CP-D/632**

**Date:** 2 April 2010  
**To:** Distribution  
**From:** N. Otsuka, O. Schwerer

**Subject:** **Resonance parameters in CPND entries**

We have been accumulating resonance parameters of compound nucleus formed in neutron-induced reactions in EXFOR. These are usually derived from neutron-induced reactions. In addition, we often see resonance parameters derived from measurement of charged-particle induced reactions. They can be divided to the following two types:

- (1) Resonance parameters of compound nucleus
- (2) Resonance parameters of residual nucleus

Below are examples of these cases in recent publications:

**(1) Resonance parameters of compound nucleus – EXFOR E2133.003 [1]**

Authors derived resonance parameters ( $E_p$  and  $\Gamma_p$ ) of  $^8\text{B}$  from measurement of proton angular distribution in  $\text{p} + ^7\text{Be} \rightarrow ^8\text{B} \rightarrow \text{p} + ^7\text{Be}$  elastic scattering in the inverse kinematics technique. These values can be easily compiled in the EXFOR format as follows

Resonance energy: (1-H-1(4-BE-7,0),,EN)  
Elastic width: (1-H-1(4-BE-7,EL),,WID)

**(2) Resonance parameters of residual nucleus – EXFOR C1707.002 [2]**

Authors derived resonance parameters ( $E_x$ ,  $\Gamma_a$ ,  $\omega\gamma$ ) of  $^{19}\text{Ne}$  from measurement of  $\alpha$  decay branching ratios of  $^{19}\text{Ne}$  populated by  $^{19}\text{F}(^3\text{He},t)^{19}\text{Ne}$  and known lifetimes for various  $^{19}\text{Ne}$  resonance states. Because they are tabulated as a function of the excitation level energy of  $^{19}\text{Ne}$ , we cannot compile these parameters as we have done for “classical” resonance parameters for neutron-induced reaction. Also we do not know what should be coded in SF1 (target) and SF2 (projectile) which forms the compound  $^{19}\text{Ne}$  resonances.

Resonance properties derived by both methods have been also used for the nuclear structure data evaluation. The XUNDL database (Experimental Unevaluated Nuclear Data List) compiled by McMaster University, Argonne National Laboratory and Triangle Universities is an example of experimental resonance parameter collections for structure data evaluators and above two data sets are compiled in XUNDL.

We propose the following restriction:

*“Resonance parameters are compiled when both the projectile and target leading to the compound resonance are clarified by authors, and also the parameters are given as a function of incident energy on resonance (compiled under data heading*

*EN-RES or, when determined in the same experiment, as REACTION with SF6=EN).*"

Note that compilers ask authors when the reference frame (lab. or c.m.) is not clear for the resonance parameters.

### **Erroneous subentries**

In this occasion, we searched some suspicious resonance parameter compilation in EXFOR. Centres are requested to correct the affected entries or, if they fall under the above restriction, to delete the affected (sub)entries.

#### **1) Resonance energy (SF6=EN) coded with SF3 ≠ 0**

13197.005  
23017.006  
31475.002-004, 31668.003  
40894.002, 40894.004, 41398.002-003, 41399.002  
C0115.006, C1502.002  
D0544.004-005  
K2002.003, K2061.003  
T0089.004, T0236.010, T0268.003-004

#### **2) Resonance spin J (SF6=J) coded with SF3 ≠ 0**

22318.004, 23017.006  
40894.002, 40894.004  
C1674.002  
E1494.006, E1494.011-012  
F0948.003  
T0236.010, T0268.003-004

#### **3) Resonance spin L (SF6=L) coded with SF3 ≠ 0**

C0295.003-004, C1561.002-004, C1634.002  
E1494.006, E1494.011-012

#### **4) Resonance parity (SF6=PTY) coded with SF3 ≠ 0**

41267.005  
C1487.002, C1634.002, C1674.002  
T0236.010, F0516.002, F0948.003  
T0268.003-004

#### **5) Resonance width (SF6=WID) coded with SF3 = 0.**

12923.003  
40383.007  
D5032.003-004, D5032.006-007, D5035.003-006  
F0632.005, F0632.007-008, F0800.007-008  
O0810.002, O1068.002, O1108.002

### **References**

- [1] H. Yamaguchi *et al*, Phys. Lett. B **672**(2009)230 (XUNDL 2009YA01)
- [2] W. P. Tan *et al.*, Phys. Rev. C **79**(2009)055805 (XUNDL 2009TA09)

**Distribution:**

blokhin@ippe.ru  
chiba@earth.sgu.ac.jp  
claes.nordborg@oecd.org  
emmeric.dupont@oecd.org  
ganesan@barc.gov.in  
gezg@ciae.ac.cn  
hongwei@ciae.ac.cn  
jhchang@kaeri.re.kr  
j.roberts@iaea.org  
kaltchenko@kinr.kiev.ua  
katakura.junichi@jaea.go.jp  
kato@nucl.sci.hokudai.ac.jp  
kiralyb@atomki.hu  
l.vrapcenjak@iaea.org  
manuel.bossant@oecd.org  
manokhin@ippe.ru  
mmarina@ippe.ru  
mwberman@bnl.gov  
nicolas.soppera@oecd.org  
nklimova@kinr.kiev.ua

n.otsuka@iaea.org  
nrdc@jcprg.org  
oblozinsky@bnl.gov  
ogritzay@kinr.kiev.ua  
otto.schwerer@aon.at  
pronyaev@ippe.ru  
r.forrest@iaea.org  
samaev@obninsk.ru  
s.babykina@polyn.kiae.su  
scyang@kaeri.re.kr  
s.dunaeva@iaea.org  
stakacs@atomki.hu  
stanislav.hlavac@savba.sk  
taova@expd.vniief.ru  
tarkanyi@atomki.hu  
varlamov@depni.sinp.msu.ru  
vlasov@kinr.kiev.ua  
vmclane@optonline.net  
v.zerkin@iaea.org  
yolee@kaeri.re.kr

**cc:**

hispin@univmail.cis.mcmaster.ca