

**NATIONAL NUCLEAR DATA CENTER
Bldg. 197D
Brookhaven National Laboratory
P. O. Box 5000
Upton, NY 11973-5000 U.S.A.**

(Internet) "NNDC@BNL.GOV"

Telephone: (516)344-2902
FAX: (516)344-2806

Memo CP-C/290

DATE: July 23, 2001
TO: Distribution
FROM: V. McLane
SUBJECT: Momentum distribution data

I have received some data from a study of the longitudinal momentum distribution using 8B beams on Pb and Ag, and propose we add a new code DP to Dictionary 32 for such data. Appropriate units will also be needed.

Proposed dictionary additions follow.

Dictionary 25 (UNITS)

MB/MEV/C	Millibarns/MeV/c	DP
----------	------------------	----

Dictionary 32 (Parameters)

DP	Differential with respect to linear momentum of outgoing particles
----	--

Dictionary 36 (Quantities)

LON, DA/DP, , IPA	Diff. with respect to longitudinal sec.mom.,int. over ang.range
-------------------	---

I have attached a sample coded entry.

Distribution:

M. Chiba, Sapporo	S. Maev, CJD
F. E. Chukreev, CAJaD	O. Schwerer, NDS
S. Dunaeva, Sarov	S. Takács, ATOMKI
O. Gritzay, KINR	F. T. Tárkányi, ATOMKI
K. Kato, JCPDG	V. Varlamov, CDFE
M. Kellett, NEADB	Zhuang Youxiang, CNDC
V. N. Manokhin, CJD	NNDC File

TRANS 20010723
 ENTRY C0820 20010723
 SUBENT C0820001 20010723
 BIB 9 17
 INSTITUTE (1USAMSU,1USAANL,2UK SUR)
 REFERENCE (J,PR/C,63,065806,2001)
 AUTHOR (B.Davids,S.M.Austin,D.Bazin,H.Esbensen,B.M.Sherrill,
 I.J.Thompson,J.A.Tostevin)
 TITLE Electromagnetic dissociation of 8B and the rate of the
 7Be(p,gamma)8B reaction in the Sun
 FACILITY (CYCLO,1USAMSU) K1200 cyclotron.
 INC-SOURCE 8B beam produced using 100 and 125 MeV/nucleon beams
 of 12C. Beam magnetically analyzed using A1200
 fragment separator.
 Beam contaminants included 7Be (5-8 times
 more intense than 8B component), 6Li, and 9C.
 ERR-ANALYS (ERR-S) Statistical uncertainty given.
 In addition, there are 9% systematic uncertainties
 due to target thickness and beam intensity.
 STATUS Data received by email from B.Davids, 26 June 2001.
 HISTORY (20010723C) VM
 ENDBIB 17 0
 NOCOMMON 0 0
 ENDSUBENT 20 0
 SUBENT C0820002 20010723
 BIB 5 14
 REACTION (47-AG-0(5-B-8,X)4-BE-7,LON,DA/DP,,IPA)
 SAMPLE 27 mg/cm2 Ag target.
 DETECTOR (MAGSP) S800 spectrometer used to detect 7Be fragments
 Spectrometer comprised of:
 . 2 position-sensitive cathode readout drift chambers
 to measure position and angles of 7Be fragments,
 . a 16 segment ionization chamber to record energy
 losses,
 . 3 plastic scintillators, the first of which measured
 total energies of particles reaching focal plane.
 METHOD (TOF) 8B particles in beam identified by time-of-
 flight.
 CORRECTION Corrected for overall efficiency of drift chambers,
 and angular acceptance of S800 spectrometer.
 ENDBIB 14 0
 COMMON 1 3
 EN
 MEV
 352.
 ENDCOMMON 3 0
 DATA 4 42

ANG-MAX	MOM-SEC	DATA	ERR-S
ADEG	MEV/C	MB/MEV/C	MB/MEV/C
1.5	1979.65	0.0806159	0.0161105
1.5	1989.0	0.296264	0.0320699
1.5	1998.35	0.497803	0.0313007
1.5	2007.7	0.642912	0.0354975
1.5	2017.05	0.693297	0.0368623
...			
2.5	2101.19	0.0665081	0.0267443

 ENDDATA 44 0
 ENDSUBENT 67 0
 ENENTRY 6 0
 ENDTRANS 1 0